CONGRESS ORTHOPAEDIC ASSOCIATES, INC.

IN MEMORIUM C.W. LAMBERT, M.D. ROSS LAMBERT, M.D. GEORGE L. MULFINGER, M.D.

ORTHOPAEDIC SURGERY
PASADENA LOCATION
RICHARD C. DIEHL, JR., M.D.
GREGORY J. ADAMSON, M.D.
JAMES A. SHANKWILER, M.D.
MICHAEL J. FRAIPONT, M.D.
WILLIAM M. COSTIGAN, M.D.
TODD B. DIETRICK, M.D.
KENNETH R. SABBAG, M.D.
THOMAS G. HARRIS, M.D.
ROY F. ASHFORD, M.D.
TIMOTHY J. JACKSON, M.D.

WWW.CONGRESSMEDICAL.COM

PASADENA 800 S. RAYMOND PASADENA, CALIFORNIA 91105 (626) 795-8051 Fax (626) 795-0356

> ARCADIA 289 W. HUNTINGTON DRIVE SUITE # 103 ARCADIA, CA 91007 (626) 821-0707 Fax (626)821-0239

P.O. BOX 90730 PASADENA, CA 91109 ORTHOPAEDIC SURGERY

ARCADIA LOCATION
GARY M. MOSCARELLO, M.D.
JOHN T. QUIGLEY, M.D.
BRADFORD H. HACK, M.D.
RISHI GARG, M.D.
JOE Y.B. LEE, M.D.
STEVEN D. LIN, M.D.
PAMELA C. LUK, M.D.

INTERNAL MEDICINE/RHEUMATOLOGY IN CHANG KIM, M.D, F.A.C.P.

PHYSICAL MEDICINE & REHABILITATION RAYMOND D. GRITTON, M.D.

OCCUPATIONAL MEDICINE WILLIAM R. DIETRICK, M.D.

ANESTHESIOLOGY AMY KIM, M.D. KALYANI V. KARANDIKAR, M.D

POST OPERATIVE INSTRUCTIONS FOR UPPER EXTREMITY SURGERY

- Keep your bandage/splint/cast clean and dry.
- 2. If you had an arthroscopic procedure, you may take your bandage off in 72 hours. There will be white tape over your incision sites. LEAVE THEM ON. At this time you may shower. Pat your elbow/wrist dry trying not to disrupt the tape and cover with a band aid. You may not soak, bathe, Jacuzzi or swim until given permission.
- 3. Elevate your upper extremity above your heart by placing it on 2 pillows with your upper body lying flat to minimize swelling and discomfort. Wiggle your fingers frequently.
- 4. You should ice your wrist/elbow off and on for the first week. If you do take your bandage off, make sure you place a cloth over the skin before placing ice back on your elbow/wrist to prevent frost bite.
- 5. Pain medication has been prescribed for you. Pain medication should be taken with food. If you are having minor discomfort, Extra Strength Tylenol will work best. Please be aware most prescription pain medications (Vicodin, Percocet, and Norco) can cause constipation. Be sure to drink plenty of water or juice.
- 6. You were given a local or regional anesthetic during surgery. This may wear off in 6 hours or longer. You may have increased pain after it has worn off.
- 7. Please call the office (626) 795-8051 (Pasadena) or (626) 821-0707 (Arcadia) to schedule your follow up appointment on ______.
- 8. If you develop a fever above 101 degrees, have increase in pain or any other concerns; please notify your surgeon's medical assistant.
- 9. If you have any further questions, please do not hesitate to call the physician's office.